

NEWS AND VIEWS

OPINION

Building strong relationships between conservation genetics and primary industry leads to mutually beneficial genomic advances

STEPHANIE J. GALLA,*
THOMAS R. BUCKLEY,†‡ ROB ELSHIRE,§
MARIE L. HALE,* MICHAEL KNAPP,¶ JOHN
MCCALLUM,** ROGER MORAGA,††
ANNA W. SANTURE,‡ PHILLIP WILCOX‡‡
and TAMMY E. STEEVES*

*School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand; †Landcare Research, Private Bag 92170, Auckland Mail Centre, Auckland 1142, New Zealand; ‡School of Biological Sciences, University of Auckland, Auckland 1010, New Zealand; §The Elshire Group, Ltd., 52 Victoria Avenue, Palmerston North 4410, New Zealand; ¶Department of Anatomy, University of Otago, P.O. Box 913, Dunedin 9054, New Zealand; **Breeding and Genomics, New Zealand Institute for Plant and Food Research, Private Bag 4704, Christchurch 8140, New Zealand; ††AgResearch, Ruakura Research Centre, Bisley Road, Private Bag 3115, Hamilton 3240, New Zealand; ‡‡Department of Mathematics and Statistics, University of Otago, P.O. Box 56, 710 Cumberland Street, Dunedin 9054, New Zealand

Several reviews in the past decade have heralded the benefits of embracing high-throughput sequencing technologies to inform conservation policy and the management of threatened species, but few have offered practical advice on how to expedite the transition from conservation genetics to conservation genomics. Here, we argue that an effective and efficient way to navigate this transition is to capitalize on emerging synergies between conservation genetics and primary industry (e.g., agriculture, fisheries, forestry and horticulture). Here, we demonstrate how building strong relationships between conservation geneticists and primary industry scientists is leading to mutually-beneficial outcomes for both disciplines. Based on our collective experience as collaborative New Zealand-based scientists, we also provide insight for forging these cross-sector relationships.

Keywords: applied research, conservation genomics, high-throughput sequencing, interdisciplinary research, next-generation sequencing

Correspondence: Stephanie J. Galla, Fax: +64 3 364 2590; E-mail: stephanie.galla@pg.canterbury.ac.nz

Received 22 April 2016; revision received 23 August 2016; accepted 24 August 2016

One does not need to read beyond the pages of *Molecular Ecology* to see how emerging technologies are revolutionizing the way we conduct research in ecology and evolutionary biology (i.e. EEB) and conservation biology. This is exemplified by rapid advances in genomics, where in the span of two decades the field of molecular ecology has grown from using Sanger technologies to sequence single target loci to using high-throughput sequencing (HTS) technologies to affordably sequence entire draft genomes (Narum *et al.* 2013; Payseur & Rieseberg 2016; Tigano & Friesen 2016). When new technologies become available, there is a tendency for reviews to be published heralding their potential to address new and exciting questions. Beyond the value of these reviews, an even more important conversation needs to take place in the peer-reviewed literature: How do we efficiently incorporate new technologies into our research repertoire to make accelerated gains in applied and fundamental science?

The field of conservation genetics is currently in transition given rapid advancements in HTS technologies. Many reviews have highlighted the promise of embracing HTS technologies in conservation (Luikart *et al.* 2003; Kohn *et al.* 2006; Primmer 2009; Allendorf *et al.* 2010; Avise 2010; Frankham 2010a; Ouborg *et al.* 2010; Angeloni *et al.* 2011; Ekblom & Galindo 2011; Funk *et al.* 2012; McCormack *et al.* 2013; Narum *et al.* 2013; Steiner *et al.* 2013; Ellegren 2014; McMahon *et al.* 2014; Shafer *et al.* 2015; Andrews *et al.* 2016; Benestan *et al.* 2016; Grueber 2016). However, as recently discussed by Shafer *et al.* (2015, 2016) and Garner *et al.* (2016), there are a limited (albeit increasing) number of published empirical studies that apply HTS data to conservation. We are aware of empirical genomic studies in EEB that are applicable to questions in conservation (e.g. Defaveri *et al.* 2013; Hoffman *et al.* 2014; Knief *et al.* 2015; Bérénos *et al.* 2016; Hess *et al.* 2016; Prince *et al.* 2016) and there are many EEB researchers applying their genomics expertise to improve conservation outcomes for threatened species, including two of our co-authors (MK, AWS). In addition to the EEB sphere, there are conservation geneticists (e.g. our co-authors SJG, TRB, MLH, TES) who are successfully venturing into conservation genomics through collaborations with colleagues in another applied discipline well versed in genomics: primary industry (a collective term referring to scientists in agriculture, fisheries, forestry and horticulture; such as our co-authors RE, JM, RM, PW). Through building these cross-sector relationships, it has become clear that there is immense potential for conservation geneticists and primary industry scientists to collaborate on applied research that addresses aligned questions using similar genomic approaches. In this opinion piece, we use our experience as a collaborative group of New

Zealand-based scientists to argue that building strong relationships between conservation genetics and primary industry can lead to improved genomic outcomes for both disciplines and offer advice on how to best build meaningful cross-sector relationships.

Conservation genetics and genomics

Before discussing mutually beneficial genomic synergies between conservation genetics and primary industry, we feel it is important to first address what conservation genetics is, what can be gained using a genomic approach and what obstacles may impede geneticists from adopting genomic technologies. Conservation genetics is a subdiscipline of conservation biology (Soulé 1985) which uses genetic data to inform the management of threatened species in collaboration with conservation practitioners (Frankham 1995, 2010b; Avise 2008; Haig *et al.* 2016). While there is overlap between the fields of conservation genetics and EEB, we distinguish conservation genetics as an applied subdiscipline with direct implications for the management of threatened species. Many threatened taxa have experienced significant population declines (i.e. demographic bottlenecks, see Keller *et al.* 1994), leading to small populations that are susceptible to genetic factors (i.e. loss of genetic diversity, inbreeding and inbreeding depression) associated with extinction risk (Frankham 1995). Conservation geneticists have traditionally used few targeted neutral genetic markers including mitochondrial sequences, microsatellites and amplified fragment length polymorphisms (AFLPs) to measure inbreeding, relatedness and genetic diversity within threatened populations, estimate population genetic structure and gene flow among threatened populations, delineate species boundaries in threatened taxa and detect hybridization and introgression between threatened and nonthreatened species (Allendorf *et al.* 2010; Ouborg *et al.* 2010).

Advancements in HTS technologies are enabling the development of genomic resources for threatened species including the *de novo* assembly and annotation of high-quality reference genomes (e.g. Li *et al.* 2014; Zhang *et al.* 2014) and characterization of a large number of genome-wide markers such as single nucleotide polymorphisms (SNPs) (e.g. Benestan *et al.* 2015; Kraus *et al.* 2015; Lemay & Russello 2015). For conservation geneticists who have traditionally used small panels of neutral genetic markers to estimate population genetic parameters above and below the species level, HTS technologies are appealing as they enable an affordable means to discover and genotype a large quantity of genome-wide SNPs (Avise 2010; McCormack *et al.* 2013; Shafer *et al.* 2015) and these large SNP data sets are more representative of genome-wide variation and can result in higher resolution estimates of population genetic parameters (Väli *et al.* 2008; Ljungqvist *et al.* 2010; Santure *et al.* 2010; Taylor *et al.* 2015). In the field of conservation genetics and EEB, a small but rapidly growing number of empirical studies have demonstrated the utility of genomic markers in estimating population genetic structure and gene flow (Bowden *et al.* 2012; Dierickx *et al.* 2015; Lew *et al.* 2015;

Oyler-McCance *et al.* 2015), estimating relatedness (Béréanos *et al.* 2016), measuring genome-wide diversity (Robinson *et al.* 2016) and detecting hybridization and introgression (Hohenlohe *et al.* 2011). We anticipate even more conservation geneticists will begin to embrace HTS technologies as empirical evidence demonstrating the superiority of using genomic markers to inform conservation decisions grows and the costs of doing so diminishes (Box 1).

The paradigm underlying many conservation genetic studies is that a genetically diverse population as measured by neutral genetic markers is also likely to be functionally diverse (Bataillon *et al.* 1996) and therefore better able to adapt to environmental change (Frankham 2005). While many have aspired to move past this paradigm, it remains entrenched in most conservation genetic studies that use neutral markers (Caballero & García-Dorado 2013; Vilas *et al.* 2015). As a result of the lack of empirical data on functional genetic diversity in species of conservation interest, beyond studies that include immunocompetence genes like those in the major histocompatibility complex and Toll-like receptors (reviewed in Grueber 2016), it has been difficult to assess the validity of this conservation genetic paradigm. Further, even if supported by empirical data, neutral genetic data might not be a suitable proxy for functional genetic data for threatened species. For example, the translocation of individuals from a large genetically diverse population to supplement a small genetically depauperate population might introduce new genetic diversity (Weeks *et al.* 2011; IUCN/SSC 2013), but it might also inadvertently lead to outbreeding depression if source and recipient populations are each locally adapted (Edmands 2007; Frankham *et al.* 2011; but see Frankham 2015; Whiteley *et al.* 2015; He *et al.* 2016).

There is exceptional interest in using a conservation genomics approach to detect regions of the genome that underlie phenotypic variation linked to fitness in threatened populations (i.e. adaptive variation; Luikart *et al.* 2003; Kohn *et al.* 2006; Ouborg *et al.* 2010; Angeloni *et al.* 2011; Harrison *et al.* 2014; Shafer *et al.* 2015). There are several methods available to study adaptive variation, including gene-mapping approaches (i.e. genome-wide association studies or GWAS, and quantitative trait loci mapping or QTL; Slate *et al.* 2010; Stapley *et al.* 2010), outlier locus analysis (Luikart *et al.* 2003; Haas & Payseur 2016) and selective sweep mapping (Pardo-Diaz *et al.* 2015). However, determining the genetic basis of phenotypic traits, especially those linked to fitness, is complex, owing to the fact that most fitness-related traits are likely to be controlled by multiple loci (Savolainen *et al.* 2013) and many are likely to be under at least some environmental influence (Falconer & Mackay 1996; Lynch & Walsh 1998). In addition, the success of these approaches is often contingent on large sample sizes (e.g. Ball 2005) which will be challenging to generate for most species of conservation concern.

While there are challenges associated with the detection of adaptive variation in threatened populations (reviewed in Shafer *et al.* 2015), there is potential to answer new questions previously not tractable by employing small sets of targeted genetic markers. In particular, an understanding

Box 1. The costs of using a conservation genomic approach. Perspectives are those of Tammy Steeves.

Since I arrived in New Zealand from Canada in 2004, I have had the privilege of developing conservation genetic management recommendations in collaboration with several Department of Conservation recovery or specialist groups to assist the recovery of endemic taonga (treasured) bird species. To date, these recommendations have been predominantly based on genetic markers, namely mitochondrial sequences or microsatellite genotypes (e.g. Steeves *et al.* 2010; Hagen *et al.* 2011; Overbeek *et al.* 2016). In collaboration with primary industry colleagues in the MapNet community (see Boxes 2 & 3), I recently assessed the direct and indirect costs associated with shifting from a conservation genetic to a conservation genomic approach and decided to develop genomic markers (SNPs) for the endangered tuturuatu/shore plover (*Thinornis novaeseelandiae*; Fig. A) and the critically endangered kakī/black stilt (*Himantopus novaeseelandiae*; Fig. B).



Fig. A Tuturuatu/Shore plover

Tuturuatu/Shore plover—I was recently invited to be an expert advisor to the Shore Plover Specialist Group. The Specialist Group was interested in sampling captive and wild birds to estimate the extent of population genetic structure and compare levels of genetic diversity, between captive and wild tuturuatu. To achieve this, I knew the cost to develop, screen and genotype ~20 polymorphic species-specific microsatellites for 94 individuals (~10K NZD) would be more than using a reduced-representation approach to simultaneously discover and genotype >20 000 SNPs for the same number of individuals (Elshire *et al.* 2011; ~8.5K NZD). I also knew it would be possible to expedite the characterization of SNPs if I was able to use a reference-guided approach. As a member of the Avian Genome Consortium, I was aware bird genomes are small, compact and highly conserved (Zhang *et al.* 2014), and that one of the newly available high-quality bird genomes (killdeer, *Charadrius vociferus*) would likely be an appropriate proxy-reference genome for SNP discovery and genotyping in tuturuatu because both

species are members of the Family Charadriidae (Card *et al.* 2014). Thus, the main driver of my decision to embrace a conservation genomic approach for tuturuatu was to ensure that I could develop a comprehensive postgraduate research project that could deliver pertinent results to the Shore Plover Specialist Group in a timely fashion.



Fig. B Kakī/Black stilt

Kakī/Black stilt – As a member of the Kakī Recovery Group, I have used species-specific genetic markers to inform the conservation genetic management of captive and wild kakī populations for many years. For example, I routinely use genetic-based measures of relatedness based on microsatellites to inform captive pairing decisions (as per Hagen *et al.* 2011). However, emerging evidence indicates genetic-based measures are relatively poor indicators of genome-wide diversity, particularly in genetically impoverished species like kakī, and a better indication of genome-wide diversity should be obtained from genomic-based measures of relatedness based on genome-wide SNPs (Taylor *et al.* 2015; Willoughby *et al.* 2015). Thus, the main driver of my decision to generate SNPs for kakī was to establish the Kakī Recovery Programme as an exemplar of 'best practice' conservation genomic management.

of the genetic basis of fitness traits will allow more robust predictions of the evolutionary potential of threatened species (Ouborg *et al.* 2010; Harrison *et al.* 2014), including a better understanding of genetic trade-offs between traits that might constrain adaptation (Slate *et al.* 2010). Further, identifying loci underlying local adaptation is likely to help identify candidate populations for conservation translocations (Seddon 2010; He *et al.* 2016). Finally, identification of

genes responsible for detrimental traits associated with inbreeding depression will have immediate impact on the management of threatened species, especially where matings between individuals are managed (e.g. captive populations; Angeloni *et al.* 2011; Harrison *et al.* 2014; Shafer *et al.* 2015).

Despite having been available for over a decade (Margulies *et al.* 2005), a limited number of publications

have applied HTS technologies to conservation (Shafer *et al.* 2015, 2016; but see Garner *et al.* 2016), with the term ‘conservation genomics gap’ first being used in 2015 to describe the paucity of conservation geneticists using HTS technologies to inform conservation management (Shafer *et al.* 2015). While there are a growing number of examples that show how genomic data are being used to inform conservation decisions (Garner *et al.* 2016; but see Shafer *et al.* 2016; see Fig. S1, Supporting information) and many conservation geneticists who are currently producing HTS data sets, there has been a substantial time lag between when these techniques have become available and uptake by the conservation research community, especially in comparison with other applied genetic disciplines like primary industry (e.g. agriculture, fisheries, forestry and horticulture; see Fig. 1). In addition, much of the uptake in conservation biology has been restricted to threatened wild fish stocks (Garner *et al.* 2016; Shafer *et al.* 2016). Of the 51 articles in Fig. 1 classified as ‘conservation genomics’, 30% pertained to the management of declining, overfished or threatened commercially fished species (e.g. Atlantic salmon, *Salmo salar*; orange roughy, *Hoplostethus atlanticus*; delta smelt, *Hypomesus transpacificus*), which provides an excellent example of how conservation genomic research can also be relevant to other applied scientific disciplines including primary industry (e.g., these articles were classified as both ‘conservation genomics’ and ‘primary industry’ in Fig. 1).

Shafer *et al.* (2015) predominantly attribute the conservation genomics gap to a persistent disconnect between academia and real-world conservation issues. We agree strong relationships between academics and conservation practitioners are crucial, but argue the conservation genomics gap as defined by Shafer *et al.* (2015) is more akin to a




Fig. 1 Number of publications using high-throughput sequencing technologies to generate genomic data in conservation (blue line) and primary industry (red line) from 2005 to 2015. Values for this graph were derived from an ISI Web of Science literature search, using inclusive terminology (see Data S1, Supporting information for details). Curve lines have been smoothed for ease of interpretation.

‘research–implementation gap’ (Knight *et al.* 2008; Hogg *et al.* 2016). Indeed, if strong relationships between academics and conservation practitioners are absent, the likelihood that *any* research will be translated into conservation action is exceptionally low (Haig *et al.* 2016). Here, we predominantly attribute the apparent shortage of conservation geneticists using HTS technologies (i.e. the conservation genomics gap *sensu stricto*) to several interconnected challenges associated with the generation, analysis and interpretation of genomic data.

Prior to identifying these interconnected challenges, we recognize some questions in conservation are still being readily addressed with genetic data (e.g. Dowling *et al.* 2015; Li *et al.* 2015a; Pacioni *et al.* 2015; Trask *et al.* 2015; Cubrinovska *et al.* 2016; Hammerly *et al.* 2016; Overbeek *et al.* 2016). We anticipate studies such as these to persist, at least in the short term, because existing panels of genetic markers remain a sufficient low-cost option in some situations (Angeloni *et al.* 2011; McCormack *et al.* 2013; McMahon *et al.* 2014). Although we acknowledge that direct cost can be a factor contributing to the conservation genomics gap, we do not think it underpins it, especially when reduced-representation approaches (e.g. restriction site-associated DNA sequencing, genotyping-by-sequencing, exome capture and RAD capture; Baird *et al.* 2008; Elshire *et al.* 2011; Jones & Good 2016; Ali *et al.* 2016) make it possible to characterize tens of thousands of SNPs in hundreds of individuals for nonmodel species at a lower cost than developing and screening relatively few novel microsatellite markers (Narum *et al.* 2013; Andrews *et al.* 2016; Box 1). Beyond direct cost, the shortage of high-quality reference genomes is an often cited impediment to SNP discovery and genotyping for nonmodel species (e.g. Allendorf *et al.* 2010; Ouborg *et al.* 2010; Shafer *et al.* 2015), particularly when approximate SNP location is of interest (e.g. Kardos *et al.* 2015). However, an ever increasing number of high-quality and high-coverage genomes are becoming available (Ellegren 2014). It has also become apparent that low-coverage draft genomes (sometimes referred to as ‘landing-pad’ or ‘skim’ genomes), or even high-quality and high-coverage genomes of closely related taxa, can enable reference-guided mapping assembly and SNP characterization in some taxa (Card *et al.* 2014; Wang *et al.* 2014). The lack of bioinformatic expertise and pipelines required to analyse large population genomic data sets has also been frequently cited as a challenge that precludes the use of HTS technologies in conservation (e.g. McCormack *et al.* 2013; Shafer *et al.* 2015). Steep analytical learning curves are generally associated with new technologies, particularly for rapidly advancing fields like genomics where bioinformatic expertise is needed to analyse large genomic data sets. However, the analysis of large population genomic data sets is no longer exceptional. For example, in regard to SNP discovery and genotyping alone, several comprehensive bioinformatic pipelines are readily available (e.g. Glaubitz *et al.* 2014; Puritz *et al.* 2014; Herten *et al.* 2015; Sovic *et al.* 2015; Melo *et al.* 2016).

Depending on the conservation genetics project at hand, one or a combination of the challenges listed above might

impede conservation geneticists from transitioning to HTS technologies. Given the recent developments in HTS technologies and the potential it has for benefitting conservation outcomes, we suggest it is time for researchers to start sharing practical advice on how to expedite the transition from conservation genetics to conservation genomics. Here, we argue that an effective and efficient way to navigate the conservation genomics gap is to capitalize on emerging synergies between conservation genetics and primary industry, and demonstrate how building strong relationships between these two disciplines is leading to mutually beneficial genomic outcomes.

Strong relationships lead to mutually beneficial genomic advances

Conservation geneticists are skilled at building strong relationships in an interdisciplinary landscape to improve conservation outcomes (Haig *et al.* 2016; Hogg *et al.* 2016). However, by pushing the boundaries of the conservation 'silo', conservation geneticists will be better able to navigate the conservation genomics gap if they forge novel relationships with scientists that have shared genomic goals, albeit in a different discipline such as primary industry (Fig. 2). As a discipline, primary industry represents a diverse group of scientists from universities, private institutions and government organizations that apply scientific data to the benefit of primary production output (e.g. meat, fish, eggs, dairy, fruits, vegetables, fibres and timber). Some of the early draft genomes were published to improve commercial outcomes, including rice (*Oryza sativa*; Goff *et al.* 2002), red jungle fowl (*Gallus gallus*; Hillier *et al.* 2004), silkworm (*Bombyx mori*; Xia *et al.* 2004) and cattle (*Bos taurus*; Schibler *et al.* 2004). With these early reference genomes and the accumulation of massive SNP data sets coupled with phenotypic data, many primary industry scientists have years of expertise with the application of genomic data. Approximately 1981 HTS studies using genomic data have been published in primary industry from 2005 to 2015, which outnumbers those produced in conservation biology by more than an order of magnitude (Fig. 1).

Conservation has already benefitted from genomic resources provided by primary industry. For example,

genomic resources developed for cattle including the draft genome (Schibler *et al.* 2004) and the bovine SNP chip (Gunderson *et al.* 2005; Steemers *et al.* 2006; Matukumalli *et al.* 2009) have been used to estimate the extent of introgression from cattle to American bison (*Bison bison*; Halbert *et al.* 2005), measure genomic variation in American and European bison (*B. bonasus*; Pertoldi *et al.* 2009) and develop genomic resources for scimitar-horned and Arabian oryx (*Oryx dammah* and *O. leucoryx*, respectively; Ogden *et al.* 2012). Similarly, genomic resources developed for domestic sheep (*Ovis aries*) have been used to describe genome-wide diversity and assess genetic rescue for bighorn sheep (*Ovis canadensis*; Poissant *et al.* 2009; Miller *et al.* 2012). Of course, there are species of mutual interest to both conservation and primary industry, including species in the fishery and forestry sectors (e.g. Monterey pine, *Pinus radiata* D. Don; New Zealand tōtara, *Podocarpus* spp.; chinook salmon, *Oncorhynchus tshawytscha*; orange roughy, *Hoplostethus atlanticus*), and therefore, genomic resources produced by one discipline can be easily used by the other (Dillon *et al.* 2013; Larson *et al.* 2014; Marshall *et al.* 2015; da Silva *et al.* 2015). We anticipate conservation geneticists may opt to use closely related commercial or model species to inform adaptive variation studies in threatened species, given that gene-mapping approaches are contingent on large sample size (Ball 2005; see discussion above) and the small census size of threatened populations may be inadequate.

Collaborations between conservation geneticists and primary industry scientists are logical because researchers in these two disciplines are beginning to address similar questions in an applied genetic discipline (see Table 1). For example, primary industry scientists have been using neutral genome-wide SNPs to calculate inbreeding coefficients in domestic sheep (Li *et al.* 2011), reconstruct parentage assignments in cattle (Hayes 2011) and calculate diversity measures for genetic improvement in poultry (red jungle fowl, Muir *et al.* 2008; domestic turkey, *Meleagris gallopavo*, Aslam *et al.* 2012). Pipelines that have been used or developed to address these questions in commercial species are likely to be of interest to conservation geneticists, but are sometimes published in discipline-specific peer-reviewed journals such as the *Journal of Dairy Science* or *Plant Biotechnology Journal* (e.g. Allen *et al.* 2012; Li *et al.* 2015b).




Fig. 2 Simplified schematic detailing how relationships between conservation genetics and primary industry are leading to mutually beneficial outcomes. In black arrows, genomic expertise from primary industry advances conservation genetics, which in turn informs conservation biology and conservation management and policy. In white arrows, biodiversity expertise informs primary industry research, which in turn improves primary production.

Table 1 Common genomic issues facing conservation genetics and parallel examples addressed by scientists in primary industry

Topic	Challenge for conservation genomics	Examples of corresponding research from primary industries
Polyploid genomes	Developing effective tools for genome-wide SNP discovery and genotyping for plants, invertebrates and some vertebrates with polyploid genomes	Genome-wide SNP studies on polyploids ¹ including wheat ² , cotton ³ , potato ⁴ and peanut ⁵
Genetic basis of adaptive variants	Discovery of variants underpinning traits of relevance to conservation including adaptive variation	Trait mapping for economically important traits using GWAS and QTL mapping ^{6,7} in rice ⁸ , dairy cattle ⁹ , pig ¹⁰ and soy bean ¹¹
Gene copy number variation	Quantifying genome-wide copy number variation and estimating its contribution to phenotypic variation	Quantifying genome-wide copy number variation and estimating its contribution to economically important traits in apple ¹² , pig ¹³ , wheat ¹⁴
Inbreeding and relatedness	Measuring inbreeding (f), detecting inbreeding depression and estimating relatedness (r) for small populations to maintain evolutionary potential	Measuring inbreeding (f), detecting inbreeding depression and estimating relatedness (r) in sheep ¹⁵ , pigs ^{16,17} and salmon ¹⁸ to enhance traits for commercial selection

¹Clevenger *et al.* (2015), ²Allen *et al.* (2012), ³Byers *et al.* (2012), ⁴Uitdewilligen *et al.* (2013), ⁵Bertioli *et al.* (2014), ⁶Collard & Mackill (2008), ⁷Hu *et al.* (2013), ⁸Begum *et al.* (2015), ⁹Li *et al.* (2015b), ¹⁰Zhang *et al.* (2015), ¹¹Zhou *et al.* (2015), ¹²Boocock *et al.* (2015), ¹³Wang *et al.* (2015), ¹⁴Wuerschum *et al.* (2015), ¹⁵Li *et al.* (2011), ¹⁶Herrero-Medrano *et al.* (2012), ¹⁷Silió *et al.* (2016), ¹⁸Dodds *et al.* (2015).

Similarly, there are some conservation genomic articles from nonacademic sources that are not represented in peer-reviewed literature (Garner *et al.* 2016). These examples highlight how relationships between conservation genetics and primary industry scientists can enable the dissemination of discipline-specific publications and will allow scientists from both disciplines to learn about recently developed pipelines.

Understanding the genetic basis of desired commercial traits is also a main focus in primary industry (Womack 2005; Tuberosa & Salvi 2006; Sellner *et al.* 2007; Collard & Mackill 2008; Neale & Kremer 2011; Sonah *et al.* 2011; Hu *et al.* 2013). Primary industry has benefitted from collaboration with researchers in human health to determine the genetic basis of phenotypic traits in complex pedigrees and structured populations using QTL mapping and GWAS (George *et al.* 2000; Aulchenko *et al.* 2007; Price *et al.* 2010). In turn, these gene-mapping approaches have been successfully applied to understanding the genetic basis of ecologically relevant traits in wild populations (Schielzeth & Husby 2014). While there are numerous research groups outside of primary industry exploring adaptive variation (e.g. Rietveld *et al.* 2013; Brachi *et al.* 2015; Chaves *et al.* 2016), we anticipate that conservation geneticists in particular will benefit from forging relationships with primary industry scientists given that both groups work in an applied discipline with species characterized by small effective population sizes. Additionally, there is potential for conservation geneticists to adopt a genomic selection approach (e.g. Hayes *et al.* 2009; Heffner *et al.* 2009) to generate breeding values to inform the selection of individuals for captive breeding. Lastly, we recognize that both conservation geneticists and primary industry researchers routinely work with species with complex genomes (Clevenger *et al.* 2015), and therefore researchers from these two disciplines have an opportunity to work together and think of creative bioinformatic solutions for species that present

bioinformatic challenges (Box 3). Given these commonalities, synergies between both conservation genetics and primary industry can lead to the development of improved HTS techniques and pipelines to address mutual problems in species of both conservation and commercial interest (Boxes 2 and 3; Table 1).

Relationships between conservation geneticists and primary industry scientists can result in improved commercial outcome for primary species as well. Conservation geneticists strive to preserve genetic diversity and the ecological and evolutionary processes that generate it (Groom *et al.* 2006; Haig *et al.* 2016). There is growing discussion among primary industry scientists regarding the need for commercial breeding programs to maximize genetic diversity and minimize inbreeding (Medugorac *et al.* 2009; Windig & Engelsma 2010; Joost *et al.* 2011; Lenstra *et al.* 2012; Pryce *et al.* 2012; Kristensen *et al.* 2015). Livestock and crops are often of a small effective population size (i.e. $N_e < 100$) due to many generations of artificial selection for desired traits and are thus susceptible to loss of genome-wide variation via inbreeding and genetic drift (Windig & Engelsma 2010; Leroy *et al.* 2013; Kristensen *et al.* 2015; Jiménez-Mena *et al.* 2016; Shepherd *et al.* 2016). There is evidence for inbreeding depression in rare breeds, such as cashmere goats (*Capra aegagrus*; Dai *et al.* 2015), Iranian Guilan sheep (Eteqadi *et al.* 2015) and Iberian pigs (*Sus scrofa*; Saura *et al.* 2015). There is also an increasing awareness of the risks associated with deploying very few genotypes, particularly in the presence of novel crop pathogens (Kim *et al.* 2015) and an increasing concern among rare breeds regarding the loss of genetic variation associated with traits that might be useful in future markets (e.g. Catalanian donkey *Equus africanus*, Gutiérrez *et al.* 2005; Famennoise poultry, Moula *et al.* 2009; black Slavonian pigs, Luković *et al.* 2012). Conservation geneticists have many years of expertise regarding the conservation genetic management strategies for threatened species (Frankham 2010a). As a result, conservation geneticists can

Box 2. Retrospective and prospective of genotyping by sequencing (GBS). Perspectives are those of Rob Elshire.

In 2007, I joined the Buckler Lab at Cornell University and the next-generation sequencing revolution simultaneously. My first task was to develop a new library preparation method for the nascent Illumina sequencing platform. The technology was not nearly as robust as it is today and the reads were very short (i.e. 32 bp in length). Our challenge was to sequence the nonrepetitive fraction of the maize (*Zea mays*) genome. To do that, we used a combination of digestion by restriction enzymes and gel-based size selection to exclude the repetitive fraction. The data generated formed the basis for the first Maize Hapmap paper (Gore *et al.* 2009). When that project neared completion, I was tasked with building a low-cost, high-throughput genotyping method as an extension of my previous work. The overall goal was to develop a genotyping system that would allow simultaneous marker discovery and genotyping and also address the issue of marker discovery bias. Other researcher groups at the time were developing similar methods, as there was a high demand for an affordable and reproducible method of genotyping and it was the next logical thing to try. One aim was to provide enough genetic markers at the right price point to enable plant breeding by genomic selection. To maximize the benefit of our work and encourage others to take what we did and create new methods appropriate for new questions, we made our work openly available. The resultant genotyping-by-sequencing (GBS) method was published in *PLoS One* in 2011 (Elshire *et al.* 2011).

We achieved our goal of developing a new genotyping method that was inexpensive, both in terms of cost per sample and cost per data point (i.e. fractions of a cent per marker). The low-cost and high-throughput nature of GBS allows plant breeders to genotype thousands of plants per cycle in genomic selection driven breeding programs (He *et al.* 2014). Primary industry programs in animal breeding have also taken up GBS. Unlike microsatellites or SNP chips, no previously generated genomic resources are necessary to deploy GBS. This allows researchers working in nonmodel species, such as orphan crops (i.e. crops of regional commercial importance, but not global), to take advantage of powerful genomic tools (Varshney *et al.* 2012). The situation for researchers in ecology and conservation biology is not dissimilar to that of those working with orphan crops. The budgets are small, resources meagre and the questions are of local importance with small (if any) obvious economic returns. It is no wonder that ecologists were among the earliest adopters of GBS.

During the development of the GBS, we tested it on species other than maize. Confident that it worked in a variety of kingdoms, we welcomed interested early adopters to the laboratory for assistance. Two of those early adopters worked in the ecology space. Dr. Thomas White worked with the invasive bank vole (*Myodes glareolus*, White *et al.* 2013) in Ireland which had small sample sizes and no reference genome. Dr. Nancy Chen studied the Florida scrub jay (*Aphelocoma coerulescens*) and developed a method using GBS data and Mendelian inheritance to improve SNP discovery (Chen *et al.* 2014). It became clear that we had developed a generally useful genomics research tool and it could be used by researchers across disciplines. We had already published the method in an open-access journal and provided analysis software under a free software licence. To allow researchers to more easily use this technology, we set up a GBS service at Cornell. By early 2016 the Cornell service had performed GBS analysis on over 1500 species.

After our initial GBS publication, a plethora of method modifications and additional software tools have emerged. The recently published epiGBS method (van Gurp *et al.* 2016) allows the interrogation of the methylome and does not require a reference genome, thereby extending the utility of the base method greatly. The GBSX toolkit (Herten *et al.* 2015) is a set of software designed to assist in the design of GBS based experiments. Many software packages have been developed to analyse GBS data (e.g. TASSEL-UNEAK, STACKS, GBS-SNP-CROPS, GLBPS; Lu *et al.* 2013; Catchen *et al.* 2013; Hapke & Thiele 2016; Melo *et al.* 2016) that are appropriate for species without reference genomes. Extensions to the molecular method and new software tools make these types of genomics approaches more broadly accessible; however, barriers to using this technology still exist in many disciplines, including the cost of laboratory and informatics setup and reservations in transitioning to new analytical tools.

Marker technology adoption has a long tail distribution. In 2013, I gave a talk on GBS at the *Molecular Markers in Horticulture Symposium*. Perusing the poster session, I found that researchers were using every type of marker technology that I knew about: from isozymes to GBS. Why were some researchers using cutting edge technologies? Why were others using antiquated, expensive and low information content technologies? Researchers in conservation genomics are in a similar situation. Across disciplines, the biological sciences are encountering rapidly changing technologies and increasingly larger data sets. Industry service providers with expert knowledge and experience, like my small New Zealand-based company (Elshire Group, Ltd.) and many others, can help bridge the gap. By developing relationships spanning human health, primary industry and conservation, as well as actively participating in research communities like MapNet (Box 3), we can work together to expedite the adoption of genomic technologies applicable to the questions at hand, effectively, efficiently and with confidence.

Box 3. Building strong interdisciplinary relationships: MapNet and VISG. Perspectives are those of Phil Wilcox.

MapNet is a genomics collaboration that was formed in 2005 by a collective of New Zealand-based researchers from agriculture, horticulture, forestry and human medical genetics that quickly identified analytical gaps in international statistical genetics research. In response, MapNet members formed the Virtual Institute of Statistical Genetics (i.e. VISG) in 2007 and successfully obtained research funding to address these gaps. Through these synergies, methods developed for large human data sets (e.g. *CNVrd*, *CNVrd2*, *selectionTools*; Nguyen *et al.* 2013, 2014; Cadzow *et al.* 2014) have been successfully applied to apple (*Malus pumila*) data to identify genes of interest in commercial species (e.g. Boocock *et al.* 2015). Other workflows, such as the *selectionTools* pipeline developed and applied to human data sets such as the *1000Genomes* human data (Cadzow *et al.* 2014), are applicable to other outcrossed species where genetic maps are available. Recently, these relationships have also expanded to include cross-sector projects with scientists from the EEB and conservation genetics sector, who are able to provide insight into how these pipelines can be more broadly applicable to other applied genetic disciplines.

Critical for these cross-sector collaborations is effective and ethical behaviours among researchers, distributed leadership, commitment to an explicitly articulated vision, and effective resourcing for method development and testing. Ongoing cost reductions in both high-throughput sequencing and genotyping will constantly challenge data analyses. Thus, collaborations among researchers in primary industry, human medical genetics, EEB and conservation genetics are an effective option to develop and apply genomic methods in a financially limited environment.

The benefits of the above-mentioned collaborations would ensure (a) relevant data analysis tools could be produced by adding relevance and utility to primary-sector researchers proposing to develop such tools, and (b) providing a platform for more efficient utilization of resources such as laboratory spaces and analytical capabilities, further reducing costs and therefore increasing data generation capacity. Collaborating with primary-sector researchers working on closely related species would also benefit conservation genetics by improving efficiency. In some cases, the same species may be endangered within its natural range, but be of commercial value in other regions – such as *Pinus radiata* D. Don, which is widely planted as an exotic in the southern hemisphere but endangered in its natural range in Baja and northern California. An additional benefit of such collaborations is valuable experience and learnings from primary-sector colleagues regarding experiment design, data analyses and interpretation of results. The MapNet collective was formed and run at essentially no cost, by utilizing the resources of collaborating institutions and labour of those who were committed to this initiative, thus such cross-sector networks are easy to establish and operate – and often professionally rewarding for all involved.

provide this biodiversity expertise to commercial species for improved primary production (Fig. 2).

Conservation biologists and primary industry scientists also share similar goals regarding how best to mitigate the impact of climate change (Kristensen *et al.* 2015). For example, plant and animal breeders are prioritizing the selection of heat-tolerant plants (Ye *et al.* 2015) and low-emission animals (Hayes *et al.* 2013) and conservation scientists are debating a role for intentional introgression of desired phenotypic traits (e.g. heat tolerance) among locally adapted species or populations (Hamilton & Miller 2015; Kovach *et al.* 2016; Miller & Hamilton 2016). Given these shared goals, there is merit for scientists in primary industry and conservation to work together to maintain the evolutionary potential of commercial and threatened species in a changing climate.

A compelling rationale for building strong relationships between primary industry and conservation biology is that scientists in both disciplines conduct applied genetic research. Whereas primary industry scientists respond to the needs of primary industry practitioners (i.e. plant and animal breeders, farmers, fishermen and loggers), conservation scientists respond to the needs of conservation practitioners (i.e. wildlife managers and policy makers; Gordon *et al.* 2014; Haig *et al.* 2016). Considering the research–

implementation gap that has been discussed in conservation genetic and genomic literature (Knight *et al.* 2008; Laikre *et al.* 2010; Shafer *et al.* 2015; Taylor & Soanes 2016), researchers from conservation genetics and primary industry can collaborate on how to best communicate research needs and results between scientists and practitioners. In the policy arena, both conservation geneticists and primary industry scientists work to develop improved policy regarding the utilization and dissemination of genetic and genomic information (e.g. the *Nagoya Protocol on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits Arising from their Utilization*, <https://www.cbd.int/abs>; the *International Treaty on Plant Genetic Resources for Food and Agriculture*, <http://www.planttreaty.org>) and we anticipate that relationships between the two disciplines will allow for discussion on how to best form policy regarding the application of genomic information to threatened and commercial species.

Cross-sector collaborations will provide exciting opportunities to strategize how best to engage with stakeholders (e.g. private landowners, local governments and research-funding bodies; Jacobson & Duff 1998; Dubbeling & Merzthal 2006); but where we see an even greater opportunity for considerable gains is for conservation geneticists

and primary industry scientists to learn from one another about the importance of building meaningful partnerships with local and indigenous communities. Partnering with these communities enriches conservation and primary industry science because it creates research projects that are informed by the traditional knowledge and needs of these communities from the initial research proposal to the final report. In New Zealand, scientists and practitioners have clear directives to engage with Māori (indigenous peoples of Aotearoa/New Zealand) regarding the management of taonga (treasured) species (i.e. *Ko Aotearoa Tēnei/This is New Zealand*, conventionally known as WAI 262, <http://www.waitangitribunal.govt.nz/>) and various approaches have been developed to facilitate such engagement (Tipene-Matua & Henaghan 2007; Wilcox *et al.* 2008; Hudson *et al.* 2010). In addition, researchers are required to consult with relevant Māori tribes (iwi or hapū) when applying to receive permits for scientific research on taonga species from the Department of Conservation. New Zealand endemic species of cultural importance include threatened species (e.g. tuturuatu/shore plover and kākī/black silt; Box 1) and commercial species (e.g. kūtāi/green-lipped mussel/Greenshell™ mussel, *Perna canaliculus*), and therefore, we urge conservation genetic and primary industry scientists to collaborate on how to build productive partnerships with relevant Māori communities to develop research that is responsive to the needs and expectations of those communities. Beyond New Zealand, researchers based in any of the 92 countries around the world that are signatories to the *Nagoya Protocol on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits Arising from their Utilization* (<https://www.cbd.int/abs/>) have an opportunity to do the same. However, we argue that as global citizens, all scientists should be acting as if their country was a signatory, because as we get closer to generating population genomic data sets that include whole genomes for species of cultural importance we need to be more aware of how these genomic resources can affect and benefit local and indigenous communities.

Moving forward

While multitasking empirical research, relationships with practitioners, stakeholders and interdisciplinary partnerships can be cumbersome, we are confident that the biggest gains in both conservation genetics and primary industry will be made under this approach. Given the mutual problems that can be solved when conservation geneticists and primary industry scientists work together, we encourage scientists in both disciplines to be leaders in interdisciplinary research and we offer the following advice on how to best forge these relationships:

Get out of your silo

The first step to building successful interdisciplinary relationships is for researchers to get out of their silos and meet people with aligned research goals across disciplines.

To accomplish this task for conservation genetics and primary industry, we advocate for small (<100 people) and diverse cross-sector meetings that allow participants from academia, government agencies and private institutions to actively engage with every presentation, especially those outside of their silos. In a New Zealand context, annual meetings such as MapNet (see Box 3), the Canterbury 'Omics Symposium and the Queenstown Research Week exemplify small, diverse, cross-sector meetings that allow scientists from both conservation and primary industry to meet and expand their research networks. For larger countries, these diverse and small meetings might be more effective on a regional vs. a national level. In addition to meetings, we encourage conservation geneticists and primary industry scientists to attend genomic and networking workshops to meet people with aligned vision for genomic research, albeit in another discipline.

Practice leadership in interdisciplinary research

The second step to forging mutually beneficial partnerships between conservation and primary industry is to actively communicate with and collaborate with researchers outside of one's silo. Doing so invariably requires leadership, respect and motivation to tackle shared problems (see Table 1), generally by expanding your own research program to incorporate collaborative interdisciplinary projects between conservation and primary industry (e.g. Banks 2004; Knowler & Bradshaw 2007; Hobbs *et al.* 2008; Blank 2013; Sardinas & Kremen 2015; Box 3). Upon launching these collaborations, it is essential that leaders from both parties open an honest dialog concerning expectations, limitations and potential hindrances to interdisciplinary work such as intellectual property issues. If collaborative groups choose to develop new methods or bioinformatic pipelines, we encourage these groups to test these tools on different species representing a wide range of genomic complexities (i.e. ploidy levels, genome size and number of repetitive elements, see Table 1) so these tools are robust and widely applicable to any research study (see also Boxes 2 and 3). We also advocate for these collaborative groups to develop methods and pipelines that are open source (see Box 2), which inspires others to use and improve upon cross-disciplinary tools. Pursuing co-funding opportunities between conservation and primary industry can be an excellent means of building mutually beneficial research collaborations, especially given that most grant providers favour collaborative proposals that tackle complex problems with broad research impact (Ledford 2015; but see also Bromham *et al.* 2016). Worldwide, there are groups that are forming to tackle complex problems through an interdisciplinary approach, including the Virtual Institute of Statistical Genetics (see Box 3) and Te Pūnaha Matatini (translated to 'the meeting place of many faces', <http://www.tepunahamatatini.ac.nz/>). As leaders from conservation and primary industry initialize interdisciplinary research, we encourage the formation and utilization of these groups to facilitate the scientific process and encourage the involvement of new partners.

Promote a community of interdisciplinary research

Leaders in both the conservation and primary industry sphere can go beyond collaborating with interdisciplinary scientists to promote a culture of interdisciplinary research. To accomplish this, we encourage editorial teams at conservation and EEB journals with a broad readership like *Molecular Ecology* to periodically invite perspective articles from colleagues in primary industry. We equate this approach to the recent decision made by the editorial team at *Animal Conservation* to invite submissions from conservation practitioners so conservation academics can better understand the needs and challenges of real-world conservation (Gordon *et al.* 2014). Leaders who are organizing meetings and conferences in primary industry, conservation and genomics can strive to incorporate cross-sector talks and break down organizational silos by minimizing field-specific sessions, as proposed by Taylor & Soanes (2016) and practised by cross-sector meetings like MapNet (see Box 3). We also challenge scientists in both primary industry and conservation to become good interdisciplinary mentors to promote a culture of interdisciplinary research. This can involve mentors in conservation and primary industry promoting genomic seasonal internships or research positions to students in different silos. Not only will this encourage an interdisciplinary field, but it will also produce well-rounded and informed early-career researchers with excellent interpersonal skills and a network of colleagues to help solve shared problems.

After relationships between conservation genetics and primary industry are forged, we do not anticipate relationships will end once genomic gains are made in both disciplines. Instead, we envision these relationships will continue to grow and enable both disciplines to problem-solve and incorporate new technologies for the improvement in threatened and commercial species. With other emerging techniques being discussed and used in both conservation and primary industry, including other *-omic* techniques (e.g. transcriptomics, proteomics, metabolomics; Diz & Calvete 2016; Todd *et al.* 2016), epigenetic studies (Verhoeven *et al.* 2016) and genome editing (Johnson *et al.* 2016), we expect conservation genetics and primary industry to continue to collaborate and solve mutual problems while incorporating new technologies in applied disciplines.

We are confident that building strong interdisciplinary relationships will enable genomic advances in both conservation genetics and primary industry. However, we appreciate our colleagues in the global conservation community may be pursuing different strategies to successfully navigate the transition from genetics to genomics and we look forward to hearing about them in due course. In the meantime, our hope is that new technologies including genomics will be effectively incorporated into applied genetic disciplines like conservation and primary industry, because there is much to gain using HTS technologies to improve outcomes for the world's threatened and commercial species.

Acknowledgements

We appreciate the constructive insights from all anonymous reviewers of this article, particularly in regard to comments that helped us shape the action plan for interdisciplinary research between conservation genetics and primary industry. We thank Margaret Paterson, Mauricio Gonzales Chang and Ilna Cubrinovska for their assistance with the literature search. We are grateful to Paul Gardner for inspiring this manuscript. We acknowledge Marty Faville for his valuable discussion. Finally, we thank Leonie Heyder for the tuturuatu image.

References

- Ali OA, O'Rourke SM, Amish SJ *et al.* (2016) RAD capture (rap-ture): flexible and efficient sequence-based genotyping. *Genetics*, **202**, 389–400.
- Allen AM, Barker GL, Wilkinson P *et al.* (2012) Discovery and development of exome-based, co-dominant single nucleotide polymorphism markers in hexaploid wheat (*Triticum aestivum* L.). *Plant Biotechnology Journal*, **11**, 279–295.
- Allendorf FW, Hohenlohe PA, Luikart G (2010) Genomics and the future of conservation genetics. *Nature Reviews Genetics*, **11**, 697–709.
- Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA (2016) Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews Genetics*, **17**, 81–92.
- Angeloni F, Wagermaker N, Vergeer P, Ouborg J (2011) Genomic toolboxes for conservation biologists. *Evolutionary Applications*, **5**, 130–143.
- Aslam ML, Bastiaansen JWM, Elferink MG *et al.* (2012) Whole genome SNP discovery and analysis of genetic diversity in Turkey (*Meleagris gallopavo*). *BMC Genomics*, **13**, 391.
- Aulchenko S, de Koning D, Haley C (2007) Genomewide rapid association using mixed model and regression: a fast and simple method for genomewide pedigree-based quantitative trait loci association analysis. *Genetics*, **177**, 577–585.
- Awise JC (2008) The history, purview, and future of conservation genetics. In: *Conservation Biology: Evolution in Action* (eds Carroll SP, Fox CW), pp. 5–15. Oxford University Press, Oxford.
- Awise JC (2010) Perspective: conservation genetics enters the genomics era. *Conservation Genetics*, **11**, 665–669.
- Baird NA, Etter PD, Atwood TS *et al.* (2008) Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS ONE*, **3**, e3376.
- Ball RD (2005) Experimental designs for reliable detection of linkage disequilibrium in unstructured random population association studies. *Genetics*, **170**, 859–873.
- Banks JE (2004) Divided culture: integrating agriculture and conservation biology. *Frontiers in Ecology and the Environment*, **2**, 537–545.
- Bataillon TM, David JL, Schoen DJ (1996) Neutral genetic markers and conservation genetics: simulated germplasm collections. *Genetics*, **144**, 409–417.
- Begum H, Spindel JE, Lalusin A *et al.* (2015) Genome-wide association mapping for yield and other agronomic traits in an elite breeding population of tropical rice (*Oryza sativa*). *PLoS ONE*, **10**, e0119873.
- Benestan LM, Gosselin T, Perrier C, Sainte-Marie B, Rochette R, Bernatchez L (2015) RAD genotyping reveals fine-scale genetic structuring and provides powerful population assignment in a widely distributed marine species, the American lobster (*Homarus americanus*). *Molecular Ecology*, **24**, 3299–3315.

- Benestan LM, Ferchaud L, Hohenlohe PA *et al.* (2016) Conservation genomics of natural and managed populations: building a conceptual and practical framework. *Molecular Ecology*, **25**, 2967–2977.
- Bérénos C, Ellis PA, Pilkington JG, Pemberton JM (2016) Genomic analysis reveals depression due to both individual and maternal inbreeding in a free-living mammal population. *Molecular Ecology*, **25**, 3152–3168.
- Bertioli DJ, Ozias-Akins P, Chu Y *et al.* (2014) The use of SNP markers for linkage mapping in diploid and tetraploid peanuts. *G3*, **4**, 89–96.
- Blank PJ (2013) Northern bobwhite response to conservation reserve program habitat and landscape attributes. *The Journal of Wildlife Management*, **77**, 68–74.
- Boocock J, Chagne D, Merriman TR, Black MA (2015) The distribution and impact of common copy-number variation in the genome of the domesticated apple, *Malus × domestica* Borkh. *BMC Genomics*, **16**, 848.
- Bowden R, MacFie TS, Myers S *et al.* (2012) Genomic tools for evolution and conservation in the chimpanzee: *Pan troglodytes ellioti* is a genetically distinct population. *PLoS Genetics*, **8**, e1002504.
- Brachi B, Meyer CG, Villoutreix R *et al.* (2015) Coselected genes determine adaptive variation in herbivore resistance throughout the native range of *Arabidopsis thaliana*. *Proceedings of the National Academy of Science*, **112**, 4032–4037.
- Bromham L, Dinnage R, Hua X (2016) Interdisciplinary research has consistently lower funding success. *Nature*, **534**, 684–687.
- Byers RL, Harker DB, Yourstone SM, Maughan PJ, Udall JA (2012) Development and mapping of SNP assays in allotetraploid cotton. *Theoretical Applied Genetics*, **124**, 1201–1214.
- Caballero A, García-Dorado A (2013) Allelic diversity and its implications for the rate of adaptation. *Genetics*, **195**, 1373–1384.
- Cadzow M, Boocock J, Nguyen HT, Wilcox P, Merriman TR, Black MA (2014) A bioinformatics workflow for detecting signatures of selection in genomic data. *Frontiers in Genetics*, **5**, 293.
- Card DC, Schield DR, Reyes-Velasco J *et al.* (2014) Two low coverage bird genomes and a comparison of reference-guided versus *de novo* genome assemblies. *PLoS ONE*, **9**, e106649.
- Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool set for population genomics. *Molecular Ecology*, **22**, 3124–3140.
- Chaves JA, Cooper EA, Hendry AP *et al.* (2016) Genomic variation at the tips of the adaptive radiation of Darwin's finches. *Molecular Ecology*, doi:10.1111/mec.13743.
- Chen N, van Hout CV, Gottipati S, Clari AG (2014) Using mendelian inheritance to improve high-throughput SNP discovery. *Genetics*, **198**, 847–857.
- Clevenger J, Chararro C, Pearl SA, Ozias-Akins P, Jackson SA (2015) Single nucleotide polymorphism identification in polyploids: a review, example, and recommendations. *Molecular Plant*, **8**, 831–846.
- Collard BCY, Mackill DJ (2008) Marker-assisted selection: an approach for precision plant breeding in the twenty-first century. *Philosophical Transactions of the Royal Society B*, **363**, 557–572.
- Cubrinovska I, Massaro M, Hale ML (2016) Assessment of hybridisation between the endangered Chatham Island black robin (*Petroica traversi*) and the Chatham Island tomtit (*Petroica macrocephala chathamensis*). *Conservation Genetics*, **17**, 259–265.
- Dai S, Wang C, Wang Z *et al.* (2015) Inbreeding and its effects on fleece traits of inner Mongolia cashmere goats. *Small Ruminant Research*, **128**, 50–53.
- Defaveri J, Viitaniemi H, Leder E, Merilä J (2013) Characterizing genic and nongenic molecular markers: comparison of microsatellites and SNPs. *Molecular Ecology Resources*, **13**, 377–392.
- Dierickx EG, Shultz AJ, Sato F, Hiraoka T, Edwards SV (2015) Morphological and genomic comparisons of Hawaiian and Japanese black-footed albatrosses (*Phoebastria nigripes*) using double digest RADseq: implications for conservation. *Evolutionary Applications*, **8**, 662–678.
- Dillon SK, Nolan MF, Matter P, Gapare WJ, Bragg JG, Southerton SG (2013) Signatures of adaptation and genetic structure among the mainland populations of *Pinus radiata* (D. Don) inferred from SNP loci. *Tree Genetics & Genomes*, **9**, 1447–1463.
- Diz AP, Calvete JJ (2016) Ecological proteomics: is the field ripe for integrating proteomics into evolutionary ecology research? *Journal of Proteomics*, **135**, 1–3.
- Dodds KG, McEwan JC, Brauning R *et al.* (2015) Construction of relatedness matrices using genotyping-by-sequencing data. *BMC Genomics*, **16**, 1047.
- Dowling TE, Anderson CD, Marsh PC, Rosenberg MS (2015) Population structure in the roundtail chub (*Gila robusta* complex) of the Gila River Basin as determined by microsatellites: evolutionary and conservation implications. *PLoS ONE*, **10**, e0139832.
- Dubbeling M, Merzthal G (2006) Sustaining urban agriculture requires the involvement of multiple stakeholders. In: *Cities Farming for Future, Urban Agriculture for Green and Productive cities* (ed. van Veenhuizen R), pp. 20–51. IDRC and IIRR Publishing, Ottawa, CA.
- Edmands S (2007) Between a rock and a hard place: evaluating the relative risks of inbreeding and outbreeding depression for conservation and management. *Molecular Ecology*, **16**, 463–475.
- Eklom R, Galindo J (2011) Applications of next generation sequencing in molecular ecology of non-model organisms. *Heredity*, **107**, 1–15.
- Ellegren H (2014) Genome sequencing and population genomics in non-model organisms. *Trends in Ecology and Evolution*, **29**, 51–63.
- Elshire RJ, Glaubitz JC, Sun Q *et al.* (2011) A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS ONE*, **6**, e19379.
- Eteqadi B, Hossein-Zadeh NG, Shadparvar AA (2015) Inbreeding effects on reproductive traits in Iranian Guilan sheep. *Tropical Animal Health and Production*, **47**, 533–539.
- Falconer DS, Mackay TFC (1996) *Introduction to Quantitative Genetics*, 4th edn. Longman, Essex.
- Frankham R (1995) Conservation genetics. *Annual Review of Genetics*, **29**, 305–327.
- Frankham R (2005) Genetics and extinction. *Biological Conservation*, **126**, 131–140.
- Frankham R (2010a) Where are we in conservation genetics and where do we need to go? *Conservation Genetics*, **11**, 661–663.
- Frankham R (2010b) Challenges and opportunities of genetic approaches to biological conservation. *Biological Conservation*, **143**, 1919–1927.
- Frankham R (2015) Genetic rescue of small inbred populations: meta-analysis reveals large and consistent benefits of gene flow. *Molecular Ecology*, **24**, 2610–2618.
- Frankham R, Ballou JD, Eldridge MD *et al.* (2011) Predicting the probability of outbreeding depression. *Conservation Biology*, **25**, 265–275.
- Funk WC, McKay JK, Hohenlohe PA, Allendorf FW (2012) Harnessing genomics for delineating conservation units. *Trends in Ecology and Evolution*, **27**, 489–496.
- Garner BA, Hand BK, Amish SJ *et al.* (2016) Genomics in conservation: case studies and bridging the gap between data and application. *Trends in Ecology and Evolution*, **31**, 81–83.
- George AW, Visscher PM, Haley CS (2000) Mapping quantitative trait loci in complex pedigrees: a two-step variance component approach. *Genetics*, **156**, 2081–2092.

- Glaubitz JC, Casstevens TM, Lu F *et al.* (2014) TASSEL-GBS: a high capacity genotyping by sequencing analysis pipeline. *PLoS ONE*, **9**, e90346.
- Goff SA, Ricke D, Lan TH *et al.* (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. Japonica). *Science*, **296**, 92–100.
- Gordon IJ, Evans DM, Garnder TWJ *et al.* (2014) Enhancing communication between conservation biologists and conservation practitioners: letter from the conservation front line. *Animal Conservation*, **17**, 1–2.
- Gore MA, Chia JM, Elshire RJ *et al.* (2009) A first-generation haplotype map of maize. *Science*, **326**, 1115–1117.
- Groom MJ, Meffe GK, Carroll CR (2006). *Principles of Conservation Biology* (No. 333.9516G7). Sinauer Associates, Sunderland.
- Grueber CE (2016) Comparative genomics for biodiversity conservation. *Computational and Structural Biotechnology*, **13**, 370–375.
- Gunderson KL, Steemers FJ, Lee G, Mendoza LG, Chee MS (2005) A genome-wide scalable SNP genotyping assay using microarray technology. *Nature*, **37**, 549–554.
- van Gurp TP, Wagermaker NC, Wouters B, Vergeer P, Ouborg JN, Verhoeven KJ (2016) epiGBS: reference-free reduced representation bisulfite sequencing. *Nature Methods*, **13**, 322–324.
- Gutiérrez JP, Marmi J, Goyache F, Jordana J (2005) Pedigree information reveals moderate to high levels of inbreeding and a weak population structure in the endangered Catalanian donkey breed. *Journal of Animal Breed Genetics*, **122**, 378–386.
- Haas RJ, Payseur BA (2016) Fifteen years of genomewide scans for selection: trends, lessons, and unaddressed genetic sources of complication. *Molecular Ecology*, **25**, 5–23.
- Hagen EN, Hale ML, Maloney RF, Steeves TE (2011) Conservation genetic management of a critically endangered New Zealand endemic bird: minimizing inbreeding in the black stilt *Himantopus novaezelandiae*. *Ibis*, **153**, 556–561.
- Haig SM, Miller MP, Bellinger R, Draheim HM, Mercer DM, Mullins TD (2016) The conservation genetics juggling act: integrating genetics and ecology, science and policy. *Evolutionary Applications*, **9**, 181–195.
- Halbert ND, Ward TJ, Schnabel RD, Taylor JF, Derr JN (2005) Conservation genomics: disequilibrium mapping of domestic cattle chromosomal segments in North American bison populations. *Molecular Ecology*, **14**, 2343–2362.
- Hamilton JA, Miller JM (2015) Adaptive introgression as a resource for management and genetic conservation in a changing climate. *Conservation Biology*, **30**, 33–41.
- Hammerly SC, de la Cerda DA, Bailey J, Johnson JA (2016) A pedigree gone bad: increased offspring survival after using DNA-based relatedness to minimize inbreeding in a captive population. *Animal Conservation*, **19**, 296–303.
- Hapke A, Thiele D (2016) GlibPSs: a toolkit for fast and accurate analyses of genotyping-by-sequencing data without a reference genome. *Molecular Ecology Resources*, **16**, 979–990.
- Harrison KA, Pavlova A, Telonis-Scott M, Sunnucks P (2014) Using genomics to characterize evolutionary potential for conservation of wild populations. *Evolutionary Applications*, **7**, 1008–1025.
- Hayes BJ (2011) Technical note: efficient parentage assignment and pedigree reconstruction with dense single nucleotide polymorphism data. *Journal of Dairy Science*, **94**, 2114–2117.
- Hayes BJ, Bowman PJ, Chamberlain AJ, Goddard ME (2009) Invited review: genomic selection in dairy cattle: progress and challenges. *Journal of Dairy Science*, **92**, 433–443.
- Hayes BJ, Lewin JA, Goddard ME (2013) The future of livestock breeding: genomic selection for efficiency, reduced emissions intensity, and adaptation. *Trends in Genetics*, **29**, 206–214.
- He J, Zhao X, Laroche A, Lu Z, Liu H, Li Z (2014) Genotyping-by-sequencing (GBS), an ultimate marker-assisted selection (MAS) tool to accelerate plant breeding. *Frontiers in Plant Science*, **5**, 484.
- He X, Johansson ML, Heath DD (2016) Role of genomics and transcriptomics in selection of reintroduction source populations. *Conservation Biology*, **30**, 1010–1018.
- Heffner EL, Sorrells ME, Jannink J (2009) Genomic selection for crop improvement. *Crop Science*, **49**, 1–12.
- Herrero-Medrano JM, Megens HJ, Crooijmans RP, Abellaneda JM, Ramis G (2012) Farm-by-farm analysis of microsatellite, mtDNA, and SNP genotype data reveals inbreeding and crossbreeding as threats to the survival of a native Spanish pig breed. *Animal Genetics*, **44**, 259–266.
- Herten K, Hestand MS, Vermeesch JR, Van Houdt JKC (2015) GBSX: a toolkit for experimental design and demultiplexing genotyping by sequencing experiments. *BMC Bioinformatics*, **16**, 73.
- Hess JE, Zendt JS, Matala AR, Narum SR (2016) Genetic basis of adult migration timing in anadromous steelhead discovered through multivariate association testing. *Proceedings of the Royal Society B*, **283**, doi:10.1098/rspb.2015/3064.
- Hillier L, Miller W, Birney E *et al.* (2004) Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature*, **432**, 695–716.
- Hobbs PR, Sayre K, Gupta R (2008) The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society Biology*, **363**, 543–555.
- Hoffman JL, Simpson F, David P *et al.* (2014) High-throughput sequencing reveals inbreeding depression in a natural population. *Proceedings of the National Academy of Science*, **111**, 3775–3780.
- Hogg C, Grueber CE, Pemberton D *et al.* (2016) “Devil Tools & Tech”: a synergy of conservation research and management practices. *Conservation Letters*, doi:10.1111/conl.12221.
- Hohenlohe PA, Amish SJ, Catchen JM, Allendorf FW, Luikart G (2011) Next-generation RAD sequencing identifies thousands of SNPs for assessing hybridization between rainbow and west-slope cutthroat trout. *Molecular Ecology Resources*, **11**, 117–122.
- Hu ZL, Park CA, Wu XL, Reecy HM (2013) Animal QTLdb: an improved database tool for livestock animal QTL/association data dissemination in the post-genome era. *Nucleic Acids Research*, **41**, 871–879.
- Hudson M, Milne M, Reynolds P, Russell K, Smith B (2010) Te Ara Tika Guidelines for Māori research ethics: a framework for researchers and ethics committee members. ISBN: 978-1-877495-03-8.
- IUCN/SSC (2013) *Guidelines for Reintroductions and Other Conservation Translocations. Version 1.0*. IUCN Species Survival Commission, Gland Switzerland; Cambridge, UK.
- Jacobson SK, Duff MD (1998) Training idiot savants: the lack of human dimensions in conservation biology. *Conservation Biology*, **12**, 263–267.
- Jiménez-Mena B, Hospital F, Bataillon T (2016) Heterogeneity in effective population size and its implications in conservation genetics and animal breeding. *Conservation Genetics Resources*, **8**, 35–41.
- Johnson JA, Altwegg R, Evans DM *et al.* (2016) Is there a future for genome-editing technologies in conservation? *Animal Conservation*, **19**, 97–101.
- Jones MR, Good JM (2016) Targeted capture in evolutionary and ecological genomics. *Molecular Ecology*, **25**, 185–202.
- Joost S, Colli L, Bonin A *et al.* (2011) Promoting collaboration between livestock and wildlife conservation genetic communities. *Conservation Genetic Resources*, **3**, 785–788.
- Kardos M, Luikart G, Allendorf FW (2015) Measuring individual inbreeding in the age of genomics: marker-based measures are better than pedigrees. *Heredity*, **115**, 63–72.

- Keller LF, Arcese P, Smith JNM, Hochachka WM, Stearns SC (1994) Selection against inbred song sparrows during a natural population bottleneck. *Nature*, **372**, 356–357.
- Kim HB, Joa JH, Song KJ (2015) Current status and prospects of kiwifruit (*Actinidia chinensis*) genomics. *Journal of Plant Biotechnology*, **42**, 342–349.
- Knief U, Hemmrich-Stanisak G, Wittig M *et al.* (2015) Quantifying realized inbreeding in wild and captive animal populations. *Heredity*, **114**, 397–403.
- Knight AT, Cowling RM, Rouget M, Balmford A, Lombard AT, Campbell BM (2008) Knowing but not doing: selecting priority conservation areas and the research-implementation gap. *Conservation Biology*, **22**, 610–617.
- Knowler D, Bradshaw B (2007) Farmer's adoption of conservation agriculture: a review and synthesis of recent research. *Food Policy*, **32**, 25–48.
- Kohn MH, Murphy WJ, Ostrander EA, Wayne RK (2006) Genomics and conservation genetics. *Trends in Ecology and Evolution*, **21**, 629–637.
- Kovach RP, Luikart G, Lower WH, Boyer MC, Muhlfeld CC (2016) Risk and efficacy of human-enabled interspecific hybridization for climate-change adaptation: response to Hamilton and Miller (2016). *Conservation Biology*, **30**, 428–430.
- Kraus RH, Vonholdt B, Cocchiara B *et al.* (2015) A single-nucleotide polymorphism-based approach for rapid and cost-effective genetic wolf monitoring in Europe based on noninvasively collected samples. *Molecular Ecology Resources*, **15**, 295–305.
- Kristensen TN, Hoffmann AA, Pertoldi C, Stronen AV (2015) What can livestock breeders learn from conservation genetics and vice versa? *Frontiers in Genetics*, **6**, 1–12.
- Laikre L, Allendorf FW, Aroner LC *et al.* (2010) Neglect of genetic diversity in implementation of the convention on biological diversity. *Conservation Biology*, **24**, 86–88.
- Larson WA, Seeb LW, Everett MV, Waples RK, Templin WD, Seeb JE (2014) Genotyping by sequencing resolves shallow population structure to inform conservation of chinook salmon (*Oncorhynchus tshawytscha*). *Evolutionary Applications*, **7**, 355–369.
- Ledford H (2015) Team science. *Nature*, **525**, 308–311.
- Lemay MA, Russello MA (2015) Genetic evidence for ecological divergence in kokanee salmon. *Molecular Ecology*, **24**, 798–811.
- Lenstra JA, Groeneveld LF, Eding J *et al.* (2012) Molecular tools and analytical approaches for the characterization of farm animal genetic diversity. *Animal Genetics*, **43**, 483–502.
- Leroy G, Mary-Huard T, Danvy S, Charvolin E, Danchin-Burge C (2013) Methods to estimate effective population size using pedigree data: examples in dog, sheep, cattle, and horse. *Genetics Selection Evolution*, **45**, 1.
- Lew RM, Finger AJ, Baerwald MR, Goodbla A, May B, Meek MH (2015) Using next-generation sequencing to assist a conservation hatchery: a single-nucleotide polymorphism panel for the genetic management of endangered delta smelt. *Transactions of the American Fisheries Society*, **144**, 767–779.
- Li M, Strandén I, Tiirikka T, Sevón-Aimonen M, Kantanen J (2011) A comparison of approaches to estimate the inbreeding coefficient and pairwise relatedness using genomic and pedigree data in a sheep population. *PLoS ONE*, **6**, e26256.
- Li S, Li B, Cheng C *et al.* (2014) Genomic signatures of near-extinction and rebirth of the crested ibis and other endangered bird species. *Genome Biology*, **15**, 557.
- Li Y, Lancaster ML, Cooper SJB, Taylor AC, Carthew SM (2015a) Population structure and gene flow in the endangered southern brown bandicoot (*Isodon obesulus obesulus*) across a fragmented landscape. *Conservation Genetics*, **16**, 331–345.
- Li X, Buitenhuis AJ, Lund MS *et al.* (2015b) Joint genome-wide association study for milk fatty acid traits in Chinese and Danish Holstein populations. *Journal of Dairy Science*, **98**, 8152–8163.
- Ljungqvist M, Åkesson M, Hansson B (2010) Do microsatellites reflect genome-wide genetic diversity in natural populations? A comment on Väli *et al.* (2008). *Molecular Ecology*, **19**, 851–855.
- Lu F, Lipka AE, Glaubitz J *et al.* (2013) Switchgrass genomic diversity, ploidy, and evolution: novel insights from a network-based SNP discovery protocol. *PLoS Genetics*, **9**, e1003215.
- Luikart G, England PR, Tallmon D, Jordan S, Taberlet P (2003) The power and promise of population genomics: from genotyping to genome typing. *Nature Reviews Genetics*, **4**, 981–994.
- Luković Z, Karolyi D, Klišanić V, Mahnet Ž, Škorput D (2012) Genetic parameters and trends for litter size in Black Slavonian pigs. 7th International Symposium on the Mediterranean Pig, ISBN: 2-85352-488-4.
- Lynch M, Walsh B (1998) *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, Sunderland.
- Margulies M, Egholm M, Altman WE *et al.* (2005) Genome sequencing in microfabricated high-density picolitre reactors. *Nature*, **437**, 376–380.
- Marshall C, Chagné D, Deusch O *et al.* (2015) A DNA-based diagnostic for differentiating among New Zealand endemic *Podocarpus*. *Tree Genetics & Genomes*, **11**, 1–13.
- Matukumalli LK, Lawley CT, Schnabel RD *et al.* (2009) Development and characterization of a high density SNP genotyping assay for cattle. *PLoS ONE*, **4**, e5350.
- McCormack JE, Hird SM, Zellmer AJ, Carstens BC, Brumfield RT (2013) Applications of next-generation sequencing to phylogeography and phylogenetics. *Molecular Phylogenetics and Evolution*, **66**, 526–538.
- McMahon BJ, Teeling EC, Höglund J (2014) How and why should we implement genomics into conservation? *Evolutionary Applications*, **7**, 999–1007.
- Medugorac I, Medugorac A, Russ I *et al.* (2009) Genetic diversity of European cattle breeds highlights the conservation value of traditional unselected breeds with high effective population size. *Molecular Ecology*, **18**, 3394–3410.
- Melo ATO, Bartaula R, Hale I (2016) GBS-SNP-CROP: a reference-optional pipeline for SNP discovery and plant germplasm characterization using variable length, paired-end genotyping-by-sequencing data. *BMC Bioinformatics*, **17**, 29.
- Miller JM, Hamilton JA (2016) Interspecies hybridization in the conservation toolbox: response to Kovach *et al.* 2016. *Conservation Biology*, **30**, 431–433.
- Miller JM, Poissant J, Hogg JT, Coltman DW (2012) Genomic consequences of genetic rescue in an insular population of bighorn sheep (*Ovis canadensis*). *Molecular Ecology*, **21**, 1583–1596.
- Moula N, Antoine-Moussiaux N, Farnir F, Leroy P (2009) Evaluation of the production performances of an endangered local poultry breed, the famennoise. *International Journal of Poultry Science*, **8**, 389–396.
- Muir WM, Wong GKS, Zhang Y *et al.* (2008) Genome-wide assessment of worldwide chicken SNP genetic diversity indicates significant absence of rare alleles in commercial breeds. *Proceedings of the National Academy of Science*, **105**, 17312–17317.
- Narum SR, Buerkle CA, Davey JW, Miller MR, Hohenlohe PA (2013) Genotyping-by-sequencing in ecological and conservation genomics. *Molecular Ecology*, **22**, 2841–2847.
- Neale DB, Kremer A (2011) Forest tree genomics: growing resources and applications. *Nature Reviews Genetics*, **12**, 111–112.
- Nguyen HT, Merrimen TR, Black MA (2013) CNVrd, a read-depth algorithm for assigning copy-number at the FCGR locus:

- population-specific tagging of copy number variation at *FCGR3B*. *PLoS ONE*, **8**, e63219.
- Nguyen HT, Merrimen TR, Black MA (2014) The CNVrd2 package: measurement of copy number at complex loci using high-throughput sequencing data. *Frontiers in Genetics*, **5**, 248.
- Ogden R, Baird J, Senn H, McEwing R (2012) The use of cross-species genome-wide arrays to discover SNP markers for conservation genetics: a case study from Arabian and scimitar-horned oryx. *Conservation Genetic Resources*, **4**, 471–473.
- Ouborg NJ, Pertoldi C, Loeschcke V, Bijlsma R, Hedrick PW (2010) Conservation genetics in transition to conservation genomics. *Trends in Genetics*, **26**, 177–187.
- Overbeek AL, Hauber ME, Brown E, Cleland S, Maloney RF, Steeves TE (2016) Evidence for brood parasitism in a critically endangered Charadriiform with implications for conservation. *Journal of Ornithology*, doi:10.1007/s/0336-016-1375-x.
- Oyler-McCance SJ, Cornman RS, Jones KL, Fike JA (2015) Genomic single-nucleotide polymorphisms confirm that Gunnison and greater sage-grouse are genetically well differentiated and that the bi-state population is distinct. *Condor*, **117**, 217–227.
- Pacioni C, Hunt H, Allentoft ME *et al.* (2015) Genetic diversity loss in a biodiversity hotspot: ancient DNA quantifies genetic decline and former connectivity in a critically endangered marsupial. *Molecular Ecology*, **24**, 5813–5828.
- Pardo-Diaz C, Salazar C, Jiggins CD (2015) Towards the identification of the loci of adaptive evolution. *Methods in Ecology and Evolution*, **6**, 445–464.
- Payseur BA, Rieseberg LH (2016) A genomic perspective on hybridization and speciation. *Molecular Ecology*, **25**, 2337–2360.
- Pertoldi C, Wójcik JM, Tokarska M *et al.* (2009) Genome variability in European and American bison detected using the BovineSNP50 bead chip. *Conservation Genetics*, **11**, 627–634.
- Poissant J, Shafer BA, Davis S *et al.* (2009) Genome-wide cross-amplification of domestic sheep microsatellites in bighorn sheep and mountain goats. *Molecular Ecology Resources*, **9**, 1121–1126.
- Price AL, Zaitlen NA, Reich D, Patterson N (2010) New approaches to population stratification in genome-wide association studies. *Nature Review Genetics*, **11**, 459–463.
- Primmer CR (2009) From conservation genetics to conservation genomics. *Annals of the New York Academy of Science*, **1162**, 357–368.
- Prince DJ, O'Rourke SM, Thompson TQ *et al.* (2016) The evolutionary basis of premature migration in Pacific salmon highlights the utility of genomics for informing conservation. *bioRxiv*, doi:10.1101/056853.
- Pryce JE, Hayes BJ, Goodard ME (2012) Novel strategies to minimize progeny inbreeding while maximizing genetic gain using genomic information. *Journal of Dairy Science*, **95**, 377–388.
- Puritz JB, Hollenbeck CM, Gold JR (2014) dDocent: a RAD seq, variant-calling pipeline designed for population genomics of non-model organisms. *PeerJ*, **2**, e431.
- Rietveld CA, Medland SE, Derringer J *et al.* (2013) GWAS of 126,559 individuals identifies genetic variants associated with educational attainment. *Science*, **340**, 1467–1471.
- Robinson JA, Vecchyo DO, Fan Z (2016) Genomic flatlining in the endangered island fox. *Current Biology*, **26**, 1183–1189.
- Santure AW, Stapley J, Ball AD *et al.* (2010) On the use of large marker panels to estimate inbreeding and relatedness: empirical and simulation studies of a pedigreed zebra finch population typed at 771 SNPs. *Molecular Ecology*, **19**, 1439–1451.
- Sardinas HX, Kremen C (2015) Pollination services from field-scale agricultural diversification may be context-dependent. *Agriculture, Ecosystems and Environment*, **207**, 17–25.
- Saura M, Fernández A, Varona L *et al.* (2015) Detecting inbreeding depression for reproductive traits in Iberian pigs using genome-wide data. *Genetics Selection Evolution*, **47**, 1.
- Savolainen O, Lascoux M, Merilä J (2013) Ecological genomics of local adaptation. *Nature Reviews Genetics*, **14**, 807–820.
- Schibler L, Roig A, Mahe MF *et al.* (2004) A first generation bovine BAC-based physical map. *Genetics selection evolution*, **36**, 105–122.
- Schielzeth H, Husby A (2014) Challenges and prospects in genome-wide quantitative trait loci mapping of standing genetic variation in natural populations. *Annals of the New York Academy of Sciences*, **1320**, 35–57.
- Seddon PJ (2010) From reintroduction to assisted colonization: moving along the conservation translocation spectrum. *Restoration Ecology*, **18**, 796–802.
- Sellner EM, Kim JW, McClure MC, Taylor KH, Schnabel RD, Taylor JF (2007) Board-invited review: applications of genomic information in livestock. *Journal of Animal Science*, **85**, 3148–3158.
- Shafer ABA, Wolf JBW, Alves PC *et al.* (2015) Genomics and the challenging translation into conservation practice. *Trends in Ecology and Evolution*, **30**, 78–87.
- Shafer ABA, Wolf JBW, Alves PC *et al.* (2016) Reply to Garner *et al.*. *Trends in Ecology and Evolution*, **31**, 83–84.
- Shepherd LD, de Lange PJ, Cox S, McLenachan PA, Roskrige NR, Lockhart PJ (2016) Evidence of a strong domestication bottleneck in the recently cultivated New Zealand endemic root crop, *Arthropodium cirratum* (Asparagaceae). *PLoS ONE*, **11**, e0152455.
- Silió L, Barragán C, Fernández AI, García-Casco J, Rodríguez MC (2016) Assessing effective population size, coancestry and inbreeding effects on litter size using the pedigree and SNP data in closed lines of the Iberian pig breed. *Journal of Animal Breeding and Genetics*, **133**, 145–154.
- da Silva AG, Appleyard SA, Upston J (2015) Establishing the evolutionary compatibility of potential sources of colonizers for overfished stocks: a population genomics approach. *Molecular Ecology*, **24**, 564–579.
- Slate J, Santure AW, Feulner PGD *et al.* (2010) Genome mapping in intensively studied wild vertebrate populations. *Trends in Ecology and Evolution*, **26**, 275–284.
- Sonah H, Deshmukh RK, Singh VP, Gupta DK, Singh NK, Sharma TR (2011) Genomic resources in horticultural crops: Status, utility and challenges. *Biotechnology Advances*, **29**, 199–209.
- Soulé ME (1985) What is conservation biology? A new synthetic discipline addresses the dynamics and problems of perturbed species, communities, and ecosystems. *BioScience*, **35**, 727–734.
- Sovic MG, Fries AC, Gibbs HL (2015) AftRAD: a pipeline for accurate and efficient *de novo* assembly of RADseq data. *Molecular Ecology Resources*, **15**, 1163–1171.
- Stapley J, Reger J, Feulner PGD *et al.* (2010) Adaptation genomics: the next generation. *Trends in Ecology and Evolution*, **25**, 705–712.
- Stearns FJ, Chang W, Lee G, Barker DL, Shen R, Gunderson KL (2006) Whole-genome genotyping with the single-base extension assay. *Nature Methods*, **3**, 31–33.
- Steeves TE, Maloney RF, Hale ML, Tylanakis JM, Gemmell NJ (2010) Genetic analyses reveal hybridization but no hybrid swarm in one of the world's rarest birds. *Molecular Ecology*, **19**, 5090–5100.
- Steiner CC, Putnam AS, Hoeck PEA, Ryder OA (2013) Conservation genomics of threatened animal species. *Annual Review of Animal Bioscience*, **1**, 261–281.
- Taylor HR, Soanes K (2016) Breaking out of the echo chamber: missed opportunities for genetics at conservation conferences. *Biodiversity and Conservation*, **25**, 1987.
- Taylor HR, Kardos MD, Ramstad KM, Allendorf FW (2015) Valid estimates of individual inbreeding coefficients from

- marker-based pedigrees are not feasible in wild populations with low allelic diversity. *Conservation Genetics*, **16**, 901–913.
- Tigano A, Friesen VL (2016) Genomics of local adaptation with gene flow. *Molecular Ecology*, **25**, 2144–2164.
- Tipene-Matua B, Henaghan M (2007) Establishing a Māori ethical framework for genetic research with Māori. In: *Genes, Society and the Future* (ed. Henaghan M), pp. 1–44. Human Genome Research Project, Dunedin, New Zealand.
- Todd EV, Black MA, Gemmell NJ (2016) The power and promise of RNA-seq in ecology and evolution. *Molecular Ecology*, **25**, 1224–1241.
- Trask AE, Bignal EM, McCracken DI, Monaghan P, Piernsey SB, Reid JM (2015) Evidence of the phenotypic expression of a lethal recessive allele under inbreeding in a wild population of conservation concern. *Journal of Animal Ecology*, **85**, 879–891.
- Tuberosa R, Salvi S (2006) Genomics-based approaches to improve drought tolerance of crops. *Trends in Plant Science*, **11**, 405–412.
- Uitdewilligen JGAML, Wolters AA, D'hoop BB, Borm TJA, Visser RGF, VanEck HJ (2013) A next-generation sequencing method for genotyping-by-sequencing of highly heterozygous autotetraploid potato. *PLoS*, **10**, e0141940.
- Väli Ü, Einarsson A, Waits L, Ellegren H (2008) To what extent do microsatellite markers reflect genome-wide genetic diversity in natural populations? *Molecular Ecology*, **17**, 3808–3817.
- Varshney RK, Ribaut J, Buckler ES, Tuberosa R, Rafalski JA, Langridge P (2012) Can genomics boost productivity of orphan crops? *Nature Biotechnology*, **30**, 1172–1176.
- Verhoeven KJF, vonHoldt BM, Sork VL (2016) Epigenetics in ecology and evolution: what we know and what we need to know. *Molecular Ecology*, **25**, 1631–1638.
- Vilas A, Pérez-Figueroa A, Quesada H, Caballero A (2015) Allelic diversity for neutral markers retains a higher adaptive potential for quantitative traits than expected heterozygosity. *Molecular Ecology*, **24**, 4419–4432.
- Wang B, Ekblom R, Bunikis I, Siitari H, Höglund J (2014) Whole genome sequencing of the black grouse (*Tetrao tetrix*): reference guided assembly suggests faster-Z and MHC evolution. *BMC Genomics*, **15**, 180.
- Wang L, Zu L, Liu X (2015) Copy number variation-based genome wide association study reveals additional variants contributing to meat quality in swine. *Scientific Reports*, **5**, 12535.
- Weeks AJ, Sgro CM, Young AG *et al.* (2011) Assessing the benefits and risks of translocations in changing environments: a genetic perspective. *Evolutionary Applications*, **4**, 709–725.
- White TA, Perkins SE, Heckel G, Searle JB (2013) Adaptive evolution during an ongoing range expansion: the invasive bank vole (*Myodes glareolus*) in Ireland. *Molecular Ecology*, **22**, 2971–2985.
- Whiteley AR, Fitzpatrick SW, Funk WC, Tallmon DA (2015) Genetic rescue to the rescue. *Trends in Ecology & Evolution*, **30**, 42–49.
- Wilcox PL, Charity JA, Roberts MR *et al.* (2008) A values-based process for cross-cultural dialogue between scientists and Māori. *Journal of the Royal Society of New Zealand*, **38**, 215–227.
- Willoughby JR, Fernandez NB, Lamb MC, Ivy JA, Lacy RC, DeWoody JA (2015) The impacts of inbreeding, drift and selection on genetic diversity in captive breeding populations. *Molecular Ecology*, **24**, 98–110.
- Windig JJ, Engelsma KA (2010) Perspectives of genomics for genetic conservation of livestock. *Conservation Genetics*, **11**, 635–641.
- Womack JE (2005) Advances in livestock genomics: opening the barn door. *Genome Research*, **15**, 1699–1705.
- Wuerschum T, Boeven PHG, Langer SM, Longin CFJ, Leiser W (2015) Multiply to conquer: copy number variations at Ppd-B1 and Vrn-A1 facilitate global adaptation in wheat. *BMC Genetics*, **16**, 96.
- Xia Q, Zhou S, Lu C *et al.* (2004) A draft sequence for the genome of the domesticated silkworm (*Bombyx mori*). *Science*, **306**, 1937–1940.
- Ye C, Tenorio F, Argayoso MA *et al.* (2015) Identifying and confirming quantitative trait loci associated with heat tolerance at flowering stage in different rice populations. *BMC Genetics*, **16**, 41.
- Zhang G, Li C, Li Q *et al.* (2014) Comparative genomics reveals insights into avian genome evolution and adaptation. *Science*, **346**, 1311–1320.
- Zhang C, Wang Z, Bruce H *et al.* (2015) Genome-wide association studies (GWAS) identify a QTL close to PRKAG3 affecting meat pH and colour in crossbred commercial pigs. *BMC Genetics*, **16**, 33.
- Zhou Z, Jiang Y, Wang Z *et al.* (2015) Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. *Nature Biotechnology*, **33**, 408–413.

S.J.G. and T.E.S. were the lead investigators on this research. S.J.G. and T.E.S. designed the research in collaboration with all authors, S.J.G. compiled the literature search, T.E.S., R.E., and P.W. contributed perspective boxes, and all authors wrote the manuscript and provided feedback on the reviewer's comments.

doi: 10.1111/mec.13837

Data accessibility

Text files including all literature search results presented in Fig. 1 and Fig. S1 (Supporting information) are available on Dryad: doi: 10.5061/dryad.32j55.

Supporting information

Additional supporting information may be found in the online version of this article.

Data S1 An ISI Web of Science literature search was conducted to compare the number of publications from 2005 to 2015 that have been produced in the fields of conservation biology and primary industry. It should be noted that the number of publications may be underestimated in both disciplines, given that conservation and primary industry researchers can publish their results in the grey literature (Garner *et al.* 2016).

Fig. S1 Growth in the number of genomic publications utilising high-throughput sequencing from 2005 to 2015 in the fields of primary industry (A) and conservation (B).